

INFN-NA-IV-93/33

DSF-T-93/33

New relations for two-dimensional Hermite polynomials

V.V.Dodonov

Lebedev Physics Institute, Leninsky Prospect 53, 117924 Moscow, Russia

and V.I.Man'ko

*Dipartimento di Scienze Fisiche Universita di Napoli "Federico II" and I.N.F.N., Sez. di Napoli**and Lebedev Physics Institute, Moscow*

Abstract

The effective formulas reducing the two-dimensional Hermite polynomials to the classical (one-dimensional) orthogonal polynomials are given. New one-parameter generating functions for these polynomials are derived. Asymptotical formulas for large values of indices are found. The applications to the squeezed one-mode states and to the time-dependent quantum harmonic oscillator are considered.

1 Introduction

The Hermite polynomials of several variables arise quite naturally almost in all problems relating to quantum systems described by means of multidimensional quadratic Hamiltonians: see, e.g., [1-4] and references therein. However, until now they were not widely used by other authors in the papers on quantum mechanics and quantum optics (for a few of exceptions see, e.g., [5,6]), due to the absence of simple explicit formulas, which would be convenient for the calculations and for the analysis of the relations obtained. The aim of the present paper is to give the expressions for multidimensional Hermite polynomials in terms of the well known classical orthogonal polynomials. Besides, we shall give new sum rules and generating functions, as well as asymptotic formulas for various combinations of the parameters. Although some of the results could be found in odd form in other references [1-11], we believe that bringing them together will be useful for the further applications. Just having in mind these applications, we pay a special attention to the specific sets of parameters defining the polynomials, which are typical for the quantum mechanical problems. We shall consider mainly the case of the Hermite polynomials of two variables, but some generalizations to higher dimensions will be also given.

The structure of the paper is as follows. In the next section we derive the explicit expressions for the two-dimensional Hermite polynomials in terms of the Jacobi, Gegenbauer, Legendre, and usual Hermite polynomials. The asymptotics of the two-dimensional Hermite polynomials of zero arguments are considered in Section 3. New sum rules and generating functions for the “diagonal” two-dimensional Hermite polynomials are given in Section 4, and their generalizations to higher dimensions are considered in Section 5. Section 6 is devoted to the physical applications. In that section we discuss briefly two problems: the photon statistics in squeezed mixed (thermal) quantum states, and the transition probabilities between the energy levels of a quantum oscillator with a time-dependent frequency.

2 Relations between two-dimensional Hermite polynomials and classical orthogonal polynomials of one variable

The two-dimensional Hermite polynomials $H_{nm}^{\{\mathbf{R}\}}(y_1, y_2)$ are defined by means of the generating function [12],

$$\exp\left[-\frac{1}{2}\mathbf{a}\mathbf{R}\mathbf{a} + \mathbf{a}\mathbf{R}\mathbf{y}\right] = \sum_{n,m=0}^{\infty} \frac{a_1^n a_2^m}{n!m!} H_{nm}^{\{\mathbf{R}\}}(\mathbf{y}). \quad (1)$$

Here a_1 and a_2 are arbitrary complex numbers combined into the two-dimensional vector $\mathbf{a} = (a_1, a_2)$:

$$\mathbf{a}\mathbf{R}\mathbf{a} = \sum_{i,k=0}^2 a_i R_{ik} a_k, \quad \mathbf{a}\mathbf{R}\mathbf{y} = \sum_{i,k=0}^2 a_i R_{ik} y_k,$$

and $\mathbf{R}$ is the symmetric matrix

$$\mathbf{R} = \begin{pmatrix} R_{11} & R_{12} \\ R_{12} & R_{22} \end{pmatrix}.$$

Let us begin with the case when $\mathbf{y} = 0$ (in the quantum mechanical applications it corresponds, for example, to the absence of an external force). Suppose that $R_{11} \neq 0$ and $R_{22} \neq 0$. Designating the left-hand side of eq. (1) as

$$E \equiv \exp \left(-\frac{1}{2} R_{11} a_1^2 - \frac{1}{2} R_{22} a_2^2 - R_{12} a_1 a_2 \right),$$

we expand it in the following Taylor's series,

$$E = \sum_{p=0}^{\infty} \frac{(-R_{11} a_1^2)^p}{2^p p!} \left[1 + 2 \frac{R_{12} a_2}{R_{11} a_1} + \frac{R_{22} a_2^2}{R_{11} a_1^2} \right]^p.$$

Introducing the notation

$$r = \frac{R_{12}}{\sqrt{R_{11} R_{22}}}, \quad \alpha = \sqrt{\frac{R_{22}}{R_{11}}} \frac{a_2}{a_1},$$

we can rewrite the square bracket in the right-hand side as

$$\left[1 + 2r\alpha + \alpha^2 \right] = \left[(\alpha + r)^2 - (r^2 - 1) \right] = (1 - r^2) \left[1 - (\rho + \gamma)^2 \right],$$

where

$$\rho = \frac{r}{\sqrt{r^2 - 1}}, \quad \gamma = \frac{\alpha}{\sqrt{r^2 - 1}}.$$

Developing the function $[1 - (\rho + \gamma)^2]^p$ with respect to variable γ we arrive at the expansion

$$E = \sum_{p=0}^{\infty} \sum_{k=0}^{2p} \frac{R_{11}^{p-\frac{k}{2}} R_{22}^{\frac{k}{2}}}{2^p p! k!} (r^2 - 1)^{p-\frac{k}{2}} a_1^{2p-k} a_2^k \frac{d^k}{d\rho^k} (1 - \rho^2)^p.$$

Taking into account the definition of the *associated Legendre functions* [12],

$$P_q^s(z) = \frac{(-1)^q}{2^q q!} (z^2 - 1)^{\frac{s}{2}} \frac{d^{q+s}}{dz^{q+s}} (1 - z^2)^q, \quad (2)$$

we obtain the following formula for the two-dimensional Hermite polynomial of zero arguments:

$$H_{nm}^{\{\mathbf{R}\}}(0, 0) = \mu_{mn}! (-1)^{\frac{m+n}{2}} R_{11}^{n/2} R_{22}^{m/2} (r^2 - 1)^{\frac{m+n}{4}} P_{(m+n)/2}^{|m-n|/2} \left(\frac{r}{\sqrt{r^2 - 1}} \right), \quad (3)$$

where

$$\mu_{mn} = \min(m, n),$$

and integers m, n must have the same parity (otherwise the right-hand side equals zero). We draw an attention to the definition of the associated Legendre functions: eq. (2) means

that we consider them in the *complex* domain, which *does not include* the segment $[-1, 1]$. Consequently, the definition adopted in the present paper differs from the definition used, e.g., in the theory of spherical harmonics and in Refs. [2,3,7], where a special case of eq. (3) was derived in connection with the calculations of the transition probabilities between the energy levels of a quantum oscillator with a time-dependent frequency. The reason is that in the generic case we can hardly expect the variable ρ to belong to the segment $[-1, 1]$ (although in some specific but important cases this is possible: see the last section of the paper).

For coinciding indices we get

$$H_{nn}^{\{\mathbf{R}\}}(0, 0) = n!(-\det \mathbf{R})^{\frac{n}{2}} P_n \left(\frac{-R_{12}}{\sqrt{-\det \mathbf{R}}} \right), \quad (4)$$

$P_n(z)$ being the usual Legendre polynomial. Using the relations between the associated Legendre functions $P_n^m(z)$, the Jacobi polynomials $P_k^{(\alpha, \beta)}(z)$, and the Gegenbauer polynomials $C_q^\gamma(z)$ [12,13],

$$P_n^m(z) = \frac{(n+m)!}{2^m n!} (z^2 - 1)^{\frac{m}{2}} P_{n-m}^{(m, m)}(z) = \frac{(2m)!}{2^m m!} (z^2 - 1)^{\frac{m}{2}} C_{n-m}^{m+\frac{1}{2}}(z), \quad n \geq m,$$

one can write eq. (3) in the following equivalent forms,

$$H_{nm}^{\{\mathbf{R}\}}(0, 0) = \frac{m!n!(-1)^{\frac{m+n}{2}}}{2^{\frac{|m-n|}{2}} \left(\frac{m+n}{2}\right)!} [R_{11}^n R_{22}^m (r^2 - 1)^{\mu_{mn}}]^{\frac{1}{2}} P_{\mu_{mn}}^{\left(\frac{|m-n|}{2}, \frac{|m-n|}{2}\right)} \left(\frac{r}{\sqrt{r^2 - 1}} \right), \quad (5)$$

$$H_{nm}^{\{\mathbf{R}\}}(0, 0) = \frac{\mu_{mn}! |n-m|! (-1)^{\frac{m+n}{2}}}{2^{\frac{|m-n|}{2}} \left(\frac{|m-n|}{2}\right)!} [R_{11}^n R_{22}^m (r^2 - 1)^{\mu_{mn}}]^{\frac{1}{2}} C_{\mu_{mn}}^{\frac{|m-n|+1}{2}} \left(\frac{r}{\sqrt{r^2 - 1}} \right). \quad (6)$$

These expressions hold for arbitrary values of the variable ρ .

For nonzero vector $\mathbf{y}$ function $H_{nm}^{\{\mathbf{R}\}}(y_1, y_2)$ can be written as a finite sum of products of the usual Hermite polynomials [8],

$$\begin{aligned} & \left(\frac{R_{11}^n R_{22}^m}{2^{n+m}} \right)^{-\frac{1}{2}} H_{nm}^{\{\mathbf{R}\}}(y_1, y_2) \\ &= \sum_{k=0}^{\mu_{mn}} \left(-\frac{2R_{12}}{\sqrt{R_{11}R_{22}}} \right)^k \frac{n!m!}{(n-k)!(m-k)!k!} H_{n-k} \left(\frac{\zeta_1}{\sqrt{2R_{11}}} \right) H_{m-k} \left(\frac{\zeta_2}{\sqrt{2R_{22}}} \right), \end{aligned} \quad (7)$$

where

$$\zeta_1 = R_{11}y_1 + R_{12}y_2, \quad \zeta_2 = R_{12}y_1 + R_{22}y_2. \quad (8)$$

The proof is straightforward. First we express the left-hand side of eq. (1) as the product of three exponential functions,

$$\exp \left(-\frac{1}{2} \mathbf{a} \mathbf{R} \mathbf{a} + \mathbf{a} \mathbf{R} \mathbf{y} \right) = \exp \left(-z_1^2 + 2z_1\eta_1 \right) \exp \left(-z_2^2 + 2z_2\eta_2 \right) \exp \left(-\frac{2R_{12}}{\sqrt{R_{11}R_{22}}} z_1 z_2 \right),$$

where

$$z_1 = a_1 \sqrt{\frac{R_{11}}{2}}, \quad z_2 = a_2 \sqrt{\frac{R_{22}}{2}}, \quad \eta_1 = \frac{\zeta_1}{\sqrt{2R_{11}}}, \quad \eta_2 = \frac{\zeta_2}{\sqrt{2R_{22}}}.$$

Then we expand each of three exponentials in the power series of z_1 , z_2 , and $z_1 z_2$, respectively, having in mind that the first two exponentials are the generating functions for the usual Hermite polynomials. Combining the terms with the powers z_1^{n-k} and z_2^{n-k} from the first two expansions with the terms $(z_1 z_2)^k$ arising in the expansion of the third exponential, $k = 0, 1, \dots, n$, one arrives at formula (7). Note that it remains valid even when $R_{11} = 0$ or $R_{22} = 0$, provided one takes the corresponding limit. For instance, if $R_{22} = 0$, then

$$H_{nm}^{\{\mathbf{R}\}}(y_1, y_2) = R_{12}^m \sum_{k=0}^{\mu_{mn}} \left(\frac{R_{11}}{2} \right)^{\frac{n-k}{2}} \frac{(-1)^k n! m! y_1^{m-k}}{(n-k)!(m-k)!k!} H_{n-k} \left(\frac{R_{11}y_1 + R_{12}y_2}{\sqrt{2R_{11}}} \right). \quad (9)$$

Proceeding to the limit $R_{11} = 0$ in eq. (9), we get the formula for the two-dimensional Hermite polynomial with matrix $\mathbf{R} = t\sigma_x$, where σ_x is the standard Pauli matrix,

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

It reads

$$H_{nm}^{\{t\sigma_x\}}(y_1, y_2) = t^{n+m} y_1^m y_2^n \sum_{k=0}^{\mu_{mn}} \frac{n! m! (-y_1 y_2 t)^{-k}}{(n-k)!(m-k)!k!}. \quad (10)$$

Taking into account the definition of the associated Laguerre polynomials [12],

$$L_n^\alpha(x) = \sum_{k=0}^n \binom{n+\alpha}{n-k} \frac{(-x)^k}{k!},$$

we can rewrite (10) as follows,

$$H_{nm}^{\{t\sigma_x\}}(y_1, y_2) = \mu_{mn}! t^{\nu_{mn}} (-1)^{\mu_{mn}} y_1^{\frac{(m-n+|m-n|)}{2}} y_2^{\frac{(n-m+|n-m|)}{2}} L_{\mu_{mn}}^{|m-n|}(ty_1 y_2), \quad (11)$$

where

$$\nu_{mn} = \max(m, n).$$

For coinciding indices we get

$$H_{nn}^{\{t\sigma_x\}}(y_1, y_2) = (-1)^n t^n n! L_n(ty_1 y_2). \quad (12)$$

The special cases of eqs. (11) and (12) were obtained earlier in [9,11].

Putting $\mathbf{y} = 0$ in eq. (7) we get a formula for $H_{nm}^{\{\mathbf{R}\}}(0, 0)$ which may be the most suitable for the numerical calculations:

$$H_{nm}^{\{\mathbf{R}\}}(0, 0) = \left(\frac{R_{11}^n R_{22}^m}{2^{n+m}} \right)^{\frac{1}{2}} \sum_{l=0}^{\left[\frac{\mu_{nm}}{2} \right]} \frac{(-1)^{\frac{n+m}{2}} n! m!}{l! (\mu_{nm} - 2l)! \left(\frac{|n-m|}{2} + l \right)!} (2r)^{\mu_{nm} - 2l}, \quad (13)$$

where m and n must have the same parity. For $m = n$ we have

$$H_{nn}^{\{\mathbf{R}\}}(0, 0) = \left(\frac{R_{11}R_{22}}{4} \right)^{\frac{n}{2}} \sum_{l=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(-1)^n (n!)^2}{(l!)^2 (n-2l)!} (2r)^{n-2l}. \quad (14)$$

Immediately from the generating function (1) we obtain the expression for $H_{nm}^{\{\mathbf{R}\}}(y_1, y_2)$ in terms of $H_{nm}^{\{\mathbf{R}\}}(0, 0)$ and variables ζ_1, ζ_2 defined by eq. (8):

$$H_{nm}^{\{\mathbf{R}\}}(y_1, y_2) = \sum_{l=0}^n \sum_{k=0}^m \binom{n}{l} \binom{m}{k} H_{lk}^{\{\mathbf{R}\}}(0, 0) \zeta_1^{n-l} \zeta_2^{m-k}. \quad (15)$$

3 Asymptotic formulas for $H_{nm}^{\{\mathbf{R}\}}(0, 0)$

The asymptotics of the functions $H_{nm}^{\{\mathbf{R}\}}(0, 0)$ for large values of indices can be derived, due to eq. (6), from the asymptotics of the Gegenbauer polynomials. Consider first the case when the argument of these polynomials, $r/\sqrt{r^2 - 1}$, does not belong to the interval $[-1, 1]$ (remind that $r = R_{12}/\sqrt{R_{11}R_{22}}$). Then the generalized Laplace-Heine formula (see Theorem 8.21.10 from ref. [6]) yields

$$C_n^\lambda(z) \approx \frac{\Gamma(n + \lambda)}{\Gamma(\lambda)n!} \zeta^n (1 - \zeta^{-2})^{-\lambda}, \quad n \gg 1, \quad \lambda \sim \mathcal{O}(1), \quad (16)$$

where

$$z = \frac{1}{2} (\zeta + \zeta^{-1}), \quad |\zeta| > 1. \quad (17)$$

The solution of eq. (17) with respect to ζ reads

$$\zeta = z + \epsilon \sqrt{z^2 - 1} = \frac{r + \epsilon}{\sqrt{r^2 - 1}}, \quad \epsilon = \pm 1, \quad (18)$$

where the choice of sign of ϵ is determined by the requirement $|\zeta| > 1$. Putting expression (18) into eqs. (6) and (16) we get for $n \gg 1$, $|n - m| \sim \mathcal{O}(1)$,

$$H_{nm}^{\{\mathbf{R}\}}(0, 0) \approx \frac{(-1)^{\frac{n+m}{2}} \epsilon^{-\frac{|n-m|+1}{2}}}{\sqrt{2\pi}} \Gamma\left(\frac{n+m+1}{2}\right) R_{11}^{n/2} R_{22}^{m/2} (r + \epsilon)^{\frac{n+m+1}{2}}. \quad (19)$$

We have used the identity [12]

$$\Gamma(z) \Gamma\left(z + \frac{1}{2}\right) = \sqrt{\pi} \Gamma(2z) 2^{1-2z}.$$

If parameters R_{12} and $\sqrt{R_{11}R_{22}}$ are real, then $\epsilon = \text{sign}(r)$, and eq. (19) can be rewritten as follows,

$$H_{nm}^{\{\mathbf{R}\}}(0, 0) \approx \frac{(-1)^{\theta_{nm}}}{\sqrt{2\pi}} \Gamma\left(\frac{n+m+1}{2}\right) R_{11}^{n/2} R_{22}^{m/2} \left(\frac{|R_{12}|}{\sqrt{R_{11}R_{22}}} + 1 \right)^{\frac{n+m+1}{2}}, \quad (20)$$

where

$$\theta_{nm} = \begin{cases} \frac{n-m}{2}, & \text{if } R_{12} < 0 \\ \frac{n+m}{2}, & \text{if } R_{12} > 0 \end{cases}.$$

However, eqs. (19) and (20) are not valid for small values of $|r|$, since for $R_{12} = 0$ and odd values of n and m they do not lead to the obvious relation $H_{nm}^{\{\mathbf{R}\}}(0, 0) \equiv 0$, which is an immediate consequence of definition (1). In this case we may use the asymptotic formula for the associated Legendre functions given in ref. [14],

$$P_n^{-m}(\cosh \xi) \approx \left(n + \frac{1}{2}\right)^{-m} \left(\frac{\xi}{\sinh \xi}\right)^{1/2} I_m\left(\left[n + \frac{1}{2}\right] \xi\right), \quad (21)$$

$$n \gg 1, \quad m \sim \mathcal{O}(1), \quad \text{Re} \xi > 0, \quad |\text{Im} \xi| < \pi,$$

$I_m(z)$ being the modified Bessel function (Bessel function of the third kind). Due to eq. (18),

$$\xi = \log \zeta = \log \left(\frac{r + \epsilon}{\sqrt{r^2 - 1}} \right).$$

Taking into account the relation

$$P_n^m(z) = \frac{(n+m)!}{(n-m)!} P_n^{-m}(z),$$

and a consequence of Stirling's formula,

$$\Gamma(x+y) \approx \Gamma(x)x^y, \quad x \gg 1, \quad y \sim \mathcal{O}(1), \quad (22)$$

we arrive at the expression

$$\begin{aligned} H_{nm}^{\{\mathbf{R}\}}(0, 0) &\approx \Gamma(\mathcal{N}_{nm})(-1)^{\frac{m+n}{2}} \left[\epsilon \mathcal{N}_{nm} R_{11}^n R_{22}^m (r^2 - 1)^{\mathcal{N}_{nm}} \log \left(\frac{r + \epsilon}{\sqrt{r^2 - 1}} \right) \right]^{1/2} \\ &\times I_{|n-m|/2} \left(\mathcal{N}_{nm} \log \left[\frac{r + \epsilon}{\sqrt{r^2 - 1}} \right] \right), \end{aligned} \quad (23)$$

where

$$\mathcal{N}_{nm} = \frac{n+m+1}{2}.$$

For real $r > 1$ eq. (23) turns into (20) provided one replaces the modified Bessel function by its asymptotic expression for a large argument [15],

$$I_a(z) \approx \frac{e^z}{\sqrt{2\pi z}}, \quad |z| \gg 1, \quad a \sim \mathcal{O}(1).$$

Now suppose that r is real, but $|r| < 1$. Then

$$\xi = \frac{1}{2} \log \left(\frac{1+|r|}{1-|r|} \right) - i \frac{\pi}{2} \epsilon, \quad \epsilon = \text{sign}(r),$$

and we may rewrite eq. (23) in terms of the usual Bessel functions, according to the relation

$$I_k(z) = i^{-k} J_k(iz).$$

It is sufficient to make calculations for $r > 0$, having in mind that the change of sign of r results in an extra factor $(-1)^n = (-1)^m$ before $H_{nm}^{\{\mathbf{R}\}}(0,0)$, due to eq. (3). Thus we get

$$\begin{aligned} H_{nm}^{\{\mathbf{R}\}}(0,0) &\approx (\epsilon i)^{\mu_{nm}} \Gamma(\mathcal{N}_{nm}) (-1)^{\frac{m+n}{2}} \left[\frac{1}{2} \mathcal{N}_{nm} R_{11}^n R_{22}^m (1-r^2)^{\mathcal{N}_{nm}} \right]^{1/2} \\ &\times \left[\pi + i \log \left(\frac{1+|r|}{1-|r|} \right) \right]^{1/2} J_{|n-m|/2} \left(\frac{1}{2} \mathcal{N}_{nm} \left[\pi + i \log \left(\frac{1+|r|}{1-|r|} \right) \right] \right). \end{aligned} \quad (24)$$

Taking into account the asymptotics of the Bessel function [12,15],

$$J_k(z) \approx \sqrt{\frac{2}{\pi z}} \cos \left(z - \frac{1}{2} k \pi - \frac{1}{4} \pi \right), \quad |\arg(z)| < \pi, \quad |z| \gg 1,$$

we can simplify eq. (24) as follows,

$$\begin{aligned} H_{nm}^{\{\mathbf{R}\}}(0,0) &\approx \sqrt{\frac{2}{\pi}} (\epsilon i)^{\mu_{nm}} \Gamma(\mathcal{N}_{nm}) (-1)^{\frac{m+n}{2}} \left[R_{11}^n R_{22}^m (1-r^2)^{\mathcal{N}_{nm}} \right]^{1/2} \\ &\times \cos \left(\frac{\pi}{2} \mu_{nm} + \frac{i}{2} \mathcal{N}_{nm} \log \left[\frac{1+|r|}{1-|r|} \right] \right) \\ &\equiv \frac{(-1)^{\frac{m+n}{2}}}{\epsilon^n \sqrt{2\pi}} \Gamma(\mathcal{N}_{nm}) [R_{11}^n R_{22}^m]^{1/2} \left[(1+|r|)^{\mathcal{N}_{nm}} + (-1)^n (1-|r|)^{\mathcal{N}_{nm}} \right]. \end{aligned} \quad (25)$$

If $r = 0$, this expression exactly equals zero for odd values of n and m (remind that m and n must have the same parity). Moreover, it coincides with eq. (20) provided $(1+|r|)^{\mathcal{N}_{nm}} \gg (1-|r|)^{\mathcal{N}_{nm}}$.

The exceptional case when coefficient ρ (the argument of the Gegenbauer or Jacobi polynomials) belongs to the interval $[-1, 1]$, so that $\rho = \cos \theta$, corresponds to $r = i \cot \theta$. Then we use the Hilb-like asymptotics of the Jacobi polynomial, given by Theorem 8.21.12 of Ref. [13],

$$P_q^{(\alpha, \beta)}(\cos \theta) \approx \frac{\Gamma(q + \alpha + 1)}{N^\alpha n!} \left(\sin \frac{\theta}{2} \right)^{-\alpha} \left(\cos \frac{\theta}{2} \right)^{-\beta} \left(\frac{\theta}{\sin \theta} \right)^{\frac{1}{2}} J_\alpha(N\theta),$$

where $N = n + (\alpha + \beta + 1)/2$, $\alpha > -1$, and β may be arbitrary number. Putting this expression into eq. (5) we arrive at the formula

$$H_{nm}^{\{\mathbf{R}\}}(0,0) \approx \frac{\nu_{nm}! [(-1)^{\mu_{nm}} \theta R_{11}^n R_{22}^m]^{1/2}}{(-1)^{\frac{n+m}{2}} \mathcal{N}_{nm}^{\frac{|n-m|}{2}} (\sin \theta)^{\mathcal{N}_{nm}}} J_{\frac{|n-m|}{2}}(\mathcal{N}_{nm} \theta), \quad (26)$$

which is valid for $\mathcal{N}_{nm} \gg 1$, $|n-m| \sim \mathcal{O}(1)$, $0 \leq \theta < \pi$.

Eq. (6) enables also to find the asyptotics of $H_{nm}^{\{\mathbf{R}\}}(0,0)$, when $|m-n| \gg 1$, $\mu_{nm} \sim \mathcal{O}(1)$. The definition of the Gegenbauer polynomial [12],

$$C_q^\lambda(z) = \sum_{k=0}^{\lfloor \frac{q}{2} \rfloor} \frac{(-1)^k \Gamma(\lambda + q - k)}{\Gamma(\lambda) k! (q - 2k)!} (2z)^{q-2k},$$

with account of eq. (22), yields for $\lambda \gg 1$

$$C_q^\lambda(z) \approx \sum_{k=0}^{\lfloor \frac{q}{2} \rfloor} \frac{(-1)^k \lambda^{q-k}}{k! (q - 2k)!} (2z)^{q-2k}.$$

Comparing this expression with the definition of the Hermite polynomial [12],

$$H_q(z) = q! \sum_{k=0}^{\lfloor \frac{q}{2} \rfloor} \frac{(-1)^k (2z)^{q-2k}}{k! (q - 2k)!},$$

we get the asymptotic formula for the Gegenbauer polynomial,

$$C_q^\lambda(z) \approx \frac{\lambda^{q/2}}{q!} H_q(z\sqrt{\lambda}), \quad \lambda \gg 1, \quad q \sim \mathcal{O}(1), \quad (27)$$

and its consequence for the two-dimensional Hermite polynomial of zero arguments,

$$\begin{aligned} H_{nm}^{\{\mathbf{R}\}}(0,0) &\approx \frac{(-1)^{\frac{m+n}{2}}}{\sqrt{\pi}} 2^{\frac{|n-m|}{2}} \Gamma\left(\frac{\nu_{nm} + 1}{2}\right) [R_{11}^n R_{22}^m (r^2 - 1)^{\mu_{nm}}]^{1/2} \\ &\times H_{\mu_{mn}}\left(\frac{r\sqrt{|n-m|}}{\sqrt{2(r^2 - 1)}}\right), \quad |n-m| \gg 1, \quad \mu_{nm} \sim \mathcal{O}(1). \end{aligned} \quad (28)$$

(We have used again eq. (22)). The limit cases of formula (28) for small and large values of the argument of the Hermite polynomial are quite clear.

4 One-parameter generating function and sum rules for the “diagonal” polynomials

Generating function (1) has two auxiliary parameters, a_1 and a_2 . However, for the “diagonal” polynomials, with $m = n$, a one-parameter generating function exists. To find it we choose vector $\mathbf{a}$ as follows,

$$a_1 = \sqrt{\lambda} \beta^*, \quad a_2 = \sqrt{\lambda} \beta,$$

where λ is a new complex parameter. Then multiplying both sides of eq. (1) by the factor $\exp(-\beta\beta^*)$ and calculating the gaussian integral

$$\begin{aligned} G(\lambda) &= \frac{i}{2\pi} \int \exp(-\frac{1}{2}\mathbf{a}\mathbf{R}\mathbf{a} + \mathbf{a}\mathbf{f} - \beta\beta^*) d\beta d\beta^* \\ &= \sum_{n,m=0}^{\infty} \lambda^m \frac{H_{nm}^{\{\mathbf{R}\}}(\mathbf{R}^{-1}\mathbf{f})}{n!m!} \frac{i}{2\pi} \int \beta^{*n} \beta^m e^{-\beta\beta^*} d\beta d\beta^*, \end{aligned}$$

with the account of the relation

$$\frac{i}{2\pi} \int \beta^{*n} \beta^m e^{-\beta\beta^*} d\beta d\beta^* = n! \delta_{nm},$$

we arrive at the formula ($\mathbf{I}$ means the unity matrix)

$$G(\lambda) \equiv \sum_{n=0}^{\infty} \frac{\lambda^n}{n!} H_{nn}^{\{\mathbf{R}\}}(\mathbf{R}^{-1}\mathbf{f}) = [\det(\lambda\sigma_x\mathbf{R} + \mathbf{I})]^{-\frac{1}{2}} \exp\left[\frac{\lambda}{2}\mathbf{f}(\lambda\sigma_x\mathbf{R} + \mathbf{I})^{-1}\sigma_x\mathbf{f}\right]. \quad (29)$$

The explicit form of the right-hand side of eq. (29) reads ($\mathbf{f} \equiv (f_1, f_2)$)

$$G(\lambda) = [1 + 2\lambda R_{12} - \lambda^2 \det \mathbf{R}]^{-\frac{1}{2}} \exp\left\{\frac{2\lambda f_1 f_2 - \lambda^2 (f_1^2 R_{22} + f_2^2 R_{11} - 2f_1 f_2 R_{12})}{2[1 + 2\lambda R_{12} - \lambda^2 \det \mathbf{R}]}\right\}. \quad (30)$$

If $\mathbf{f} = 0$, then this function is nothing but the generating function of the Legendre polynomials [12], and we again arrive at eq. (4).

Eq. (30) enables us to represent the “diagonal” two-dimensional Hermite polynomials in the form of a finite sum over the products of the Laguerre polynomials. For this purpose we rewrite the right-hand side of eq. (30) as follows,

$$G(\lambda) = \left[\left(1 - \frac{\lambda}{\lambda_1}\right)\left(1 - \frac{\lambda}{\lambda_2}\right)\right]^{-\frac{1}{2}} \exp\left[\frac{\lambda x_1}{\lambda - \lambda_1} + \frac{\lambda x_2}{\lambda - \lambda_2}\right],$$

where

$$\begin{aligned} \lambda_1 &= (\sqrt{R_{11}R_{22}} - R_{12})^{-1}, & \lambda_2 &= -(\sqrt{R_{11}R_{22}} + R_{12})^{-1}, \\ x_1 &= \frac{f_1^2 R_{22} + f_2^2 R_{11} - 2\sqrt{R_{11}R_{22}} f_1 f_2}{4\sqrt{R_{11}R_{22}}(\sqrt{R_{11}R_{22}} - R_{12})}, \\ x_2 &= \frac{f_1^2 R_{22} + f_2^2 R_{11} + 2\sqrt{R_{11}R_{22}} f_1 f_2}{4\sqrt{R_{11}R_{22}}(\sqrt{R_{11}R_{22}} + R_{12})}. \end{aligned}$$

Using the known relation [12] for the Laguerre polynomials

$$\frac{1}{\sqrt{1-z}} \exp \frac{xz}{z-1} = \sum_{n=0}^{\infty} z^n L_n^{-\frac{1}{2}}(x),$$

we see that the generating function $G(\lambda)$ can be represented as

$$G(\lambda) = \sum_{s=0}^{\infty} \left(\frac{\lambda}{\lambda_1} \right)^s L_s^{-\frac{1}{2}}(x_1) \sum_{k=0}^{\infty} \left(\frac{\lambda}{\lambda_2} \right)^k L_k^{-\frac{1}{2}}(x_2).$$

Thus we obtain the formula

$$H_{nn}^{\{\mathbf{R}\}}(\mathbf{R}^{-1}\mathbf{f}) = (-1)^n n! \sum_{s=0}^n (R_{12} - \sqrt{R_{11}R_{22}})^s (R_{12} + \sqrt{R_{11}R_{22}})^{n-s} L_s^{-\frac{1}{2}}(x_1) L_{n-s}^{-\frac{1}{2}}(x_2). \quad (31)$$

Assuming $\mathbf{f} = \mathbf{R}\mathbf{y}$, we get quite the same decomposition (31) for the function $H_{nn}^{\{\mathbf{R}\}}(\mathbf{y})$, but with the variables x_1 and x_2 expressed directly in terms of y_1 and y_2 as follows,

$$x_1 = \frac{1}{4} \left(1 - \frac{R_{12}}{\sqrt{R_{11}R_{22}}} \right) \left(y_1^2 R_{11} + y_2^2 R_{22} - 2\sqrt{R_{11}R_{22}} y_1 y_2 \right),$$

$$x_2 = \frac{1}{4} \left(1 + \frac{R_{12}}{\sqrt{R_{11}R_{22}}} \right) \left(y_1^2 R_{11} + y_2^2 R_{22} + 2\sqrt{R_{11}R_{22}} y_1 y_2 \right).$$

Formulas like (30) and (31) for the diagonal matrix elements of the Gaussian density matrix in the Fock basis were obtained also in Ref. [16].

Differentiating the generating function $G(\lambda)$ k times with respect to λ and putting $\lambda = 1$ in the final expression, one can calculate the cumulants $\langle n(n-1) \cdots (n-k+1) \rangle$, defined according to the relation

$$\sum_{n=0}^{\infty} n(n-1) \cdots (n-k+1) \frac{H_{nn}^{\{\mathbf{R}\}}(\mathbf{y})}{n!} = \langle n(n-1) \cdots (n-k+1) \rangle \sum_{n=0}^{\infty} \frac{H_{nn}^{\{\mathbf{R}\}}(\mathbf{y})}{n!}.$$

The first two derivatives yield the “average value” $\langle n \rangle$ and the “dispersion”

$$\sigma_n = \langle n^2 \rangle - \langle n \rangle^2 = \langle n(n-1) \rangle + \langle n \rangle - \langle n \rangle^2.$$

Differentiating the preexponential factor in eq. (29) with the aid of the matrix formula

$$\frac{d}{dt} \log \det \mathbf{A}(t) = \text{Tr} \left(\mathbf{A}^{-1} \frac{d\mathbf{A}}{dt} \right),$$

we get the following general concise relations,

$$\langle n \rangle(\mathbf{R}, \mathbf{y}) = \frac{1}{2} \text{Tr} \mathbf{S} - 1 + \frac{1}{2} \mathbf{y} \mathbf{R} \mathbf{S}^2 \sigma_x \mathbf{R} \mathbf{y}, \quad (32)$$

$$\sigma_n(\mathbf{R}, \mathbf{y}) = \frac{1}{2} \text{Tr} [\mathbf{S}(\mathbf{S} - \mathbf{I})] + \frac{1}{2} \mathbf{y} \mathbf{R} (2\mathbf{S} - \mathbf{I}) \mathbf{S}^2 \sigma_x \mathbf{R} \mathbf{y}, \quad (33)$$

where matrix $\mathbf{S}$ is defined as follows,

$$\mathbf{S} = (\sigma_x \mathbf{R} + \mathbf{I})^{-1}.$$

Of course, these formulas hold provided the series converge. The simplest special case corresponds to the matrix $\mathbf{R} = t\sigma_x$, when

$$\langle n \rangle = -\frac{t}{1+t} + y_1 y_2 \left(\frac{t}{1+t} \right)^2, \quad \sigma_n = -\frac{t}{(1+t)^2} + y_1 y_2 \frac{t^2(1-t)}{(1+t)^3}.$$

These relations have sense for $-1 < t < 0$.

5 Generalizations to higher dimensions

To generalize the results of the previous section to the “diagonal” Hermite polynomials of $2N$ variables we introduce N complex parameters λ_i , $i = 1, 2, \dots, N$, and choose the auxiliary parameters a_j as follows,

$$a_i = \sqrt{\lambda_i} \beta_i^*, \quad a_{i+N} = \sqrt{\lambda_i} \beta_i, \quad i = 1, 2, \dots, N.$$

Then the same scheme results in the following expansion,

$$\begin{aligned} & \sum_{n_1=0}^{\infty} \dots \sum_{n_N=0}^{\infty} \frac{\lambda_1^{n_1}}{n_1!} \frac{\lambda_2^{n_2}}{n_2!} \dots \frac{\lambda_N^{n_N}}{n_N!} H_{n_1 n_2 \dots n_N n_1 n_2 \dots n_N}^{\{\mathbf{R}\}}(\mathbf{y}) \\ &= [\det(\Lambda \Sigma_x \mathbf{R} + \mathbf{I}_{2N})]^{-\frac{1}{2}} \exp \left[\frac{1}{2} \mathbf{y} \mathbf{R} (\Lambda \Sigma_x \mathbf{R} + \mathbf{I}_{2N})^{-1} \Sigma_x \Lambda \mathbf{R} \mathbf{y} \right]. \end{aligned} \quad (34)$$

Here $\mathbf{y} = (y_1, y_2, \dots, y_{2N})$, the $2N \times 2N$ matrix Σ_x is the $2N$ -dimensional analog of the Pauli matrix σ_x ,

$$\Sigma_x = \begin{pmatrix} 0 & \mathbf{I}_N \\ \mathbf{I}_N & 0 \end{pmatrix},$$

$\mathbf{I}_N$ is $N \times N$ unit matrix, and the diagonal $2N \times 2N$ - matrix Λ reads

$$\Lambda = \sum_{j=1}^N \lambda_j \Lambda_j,$$

where each matrix Λ_j has only two nonzero elements:

$$(\Lambda_j)_{jj} = (\Lambda_j)_{j+N, j+N} = 1.$$

Note that matrix Λ commutes with matrix Σ_x but does not commute, in general, with matrix $\mathbf{R}$.

Generating function (34) enables to calculate the “average values” $\langle n_1^{\alpha_1} n_2^{\alpha_2} \dots n_N^{\alpha_N} \rangle$, defined as follows,

$$\begin{aligned} & \sum_{n_1=0}^{\infty} \dots \sum_{n_N=0}^{\infty} n_1^{\alpha_1} n_2^{\alpha_2} \dots n_N^{\alpha_N} \frac{H_{n_1 n_2 \dots n_N n_1 n_2 \dots n_N}^{\{\mathbf{R}\}}(\mathbf{y})}{n_1! n_2! \dots n_N!} \\ &= \langle n_1^{\alpha_1} n_2^{\alpha_2} \dots n_N^{\alpha_N} \rangle \sum_{n_1=0}^{\infty} \dots \sum_{n_N=0}^{\infty} \frac{H_{n_1 n_2 \dots n_N n_1 n_2 \dots n_N}^{\{\mathbf{R}\}}(\mathbf{y})}{n_1! n_2! \dots n_N!}. \end{aligned}$$

Taking into account the relations

$$\frac{\partial \Lambda}{\partial \lambda_j} = \Lambda_j, \quad \Lambda_j \Lambda_k = \mathbf{0}, \quad j \neq k, \quad \Lambda_j^2 = \Lambda_j, \quad \Lambda_j \Sigma_x = \Sigma_x \Lambda_j,$$

we get the following formulas,

$$\langle n_j \rangle = \frac{1}{2} \text{Tr} [\Lambda_j \mathbf{S}] - 1 + \frac{1}{2} \mathbf{y} \mathbf{R} \mathbf{S} \Lambda_j \mathbf{S} \Sigma_x \mathbf{R} \mathbf{y}, \quad (35)$$

$$\sigma_{n_j n_k} = \frac{1}{2} \text{Tr} [\Lambda_j \mathbf{S} \Lambda_k \mathbf{S}] + \frac{1}{2} \mathbf{y} \mathbf{R} \mathbf{S} \Lambda_j \mathbf{S} \Lambda_k \mathbf{S} \Sigma_x \mathbf{R} \mathbf{y}, \quad j \neq k \quad (36)$$

$$\sigma_{n_j n_j} = \frac{1}{2} \text{Tr} [\Lambda_j \mathbf{S} (\Lambda_j \mathbf{S} - \mathbf{I}_{2N})] + \frac{1}{2} \mathbf{y} \mathbf{R} (2\mathbf{S} \Lambda_j - \mathbf{I}_{2N}) \mathbf{S} \Lambda_j \mathbf{S} \Sigma_x \mathbf{R} \mathbf{y}, \quad (37)$$

where

$$\sigma_{n_j n_k} = \langle n_j n_k \rangle - \langle n_j \rangle \langle n_k \rangle,$$

and matrix $\mathbf{S}$ reads

$$\mathbf{S} = (\Sigma_x \mathbf{R} + \mathbf{I}_{2N})^{-1}.$$

6 Physical applications

1. Suppose we have some *Gaussian distribution function*

$$\mathcal{W}(\mathbf{q}) = [\det(2\pi\mathbf{M})]^{-\frac{1}{2}} \exp \left\{ -\frac{1}{2} (\mathbf{q} - \langle \mathbf{q} \rangle) \mathbf{M}^{-1} (\mathbf{q} - \langle \mathbf{q} \rangle) \right\}, \quad (38)$$

where $\mathbf{q}$ is an N -dimensional vector, and $\mathbf{M}$ is a symmetric positively definite $N \times N$ -matrix, which is nothing but the covariance matrix,

$$\mathbf{M} = \|\mathcal{M}_{ij}\|, \quad \mathcal{M}_{ij} = \langle q_i q_j \rangle - \langle q_i \rangle \langle q_j \rangle,$$

$$\langle f(\mathbf{q}) \rangle \equiv \int f(\mathbf{q}) \mathcal{W}(\mathbf{q}) d^N \mathbf{q}, \quad \int \mathcal{W}(\mathbf{q}) d^N \mathbf{q} = 1.$$

In this case the *characteristic function* $\chi(\mathbf{a}) \equiv \langle \exp(\mathbf{a} \mathbf{q}) \rangle$ is also Gaussian,

$$\chi(\mathbf{a}) = \exp \left(\frac{1}{2} \mathbf{a} \mathbf{M} \mathbf{a} + \mathbf{a} \langle \mathbf{q} \rangle \right). \quad (39)$$

On the other hand, evidently,

$$\chi(\mathbf{a}) = \sum_{n_1, \dots, n_N=0}^{\infty} \frac{a_1^{n_1} \cdots a_N^{n_N}}{n_1! \cdots n_N!} \langle q_1^{n_1} \cdots q_N^{n_N} \rangle.$$

Comparing these formulas with eq. (1) we arrive at the relation

$$\langle q_1^{n_1} \cdots q_N^{n_N} \rangle = H_{n_1 \dots n_N}^{\{-\mathbf{M}\}} (-\mathbf{M}^{-1} \langle \mathbf{q} \rangle). \quad (40)$$

2. If $N = 2$, $\mathbf{q} = (p, q)$, and $\mathcal{W}(\mathbf{q})$ is the *Wigner function* of some one-dimensional quantum system with quadrature components p, q , then the probability to observe n quanta in the state described by $\mathcal{W}(\mathbf{q})$ (in general, it is a *mixed squeezed quantum state*) is given by the relation [3,5,6,9,17]

$$\mathcal{P}_n = \mathcal{P}_0 \frac{H_{nn}^{\{\mathbf{R}\}}(y_1, y_2)}{n!}, \quad (41)$$

with the following elements of matrix $\mathbf{R}$,

$$R_{11} = R_{22}^* = \frac{2(\sigma_{pp} - \sigma_{qq} - 2i\sigma_{pq})}{1 + 2T + 4d}, \quad R_{12} = \frac{1 - 4d}{1 + 2T + 4d}, \quad (42)$$

and vector $\mathbf{y}$,

$$y_1 = y_2^* = \frac{2[(2\sigma_{pp} - 1 + 2i\sigma_{pq})\langle q \rangle + i(1 - 2\sigma_{qq} + 2i\sigma_{pq})\langle p \rangle]}{2T - 4d - 1}. \quad (43)$$

Here

$$d = \det \mathbf{M} \geq \frac{1}{4}, \quad T = \text{Tr} \mathbf{M} \geq 1 \quad (44)$$

(the inequalities are the consequences of the *generalized uncertainty relations* [3], which ensure the positive definiteness of the statistical operator corresponding to function $\mathcal{W}(\mathbf{q})$).

The probability to have no photons $\mathcal{P}_0$ is given by the formula

$$\mathcal{P}_0 = (d + \frac{1}{2}T + \frac{1}{4})^{-\frac{1}{2}} \exp \left[-\frac{\langle p \rangle^2(2\sigma_{qq} + 1) + \langle q \rangle^2(2\sigma_{pp} + 1) - 4\sigma_{pq}\langle p \rangle\langle q \rangle}{1 + 2T + 4d} \right]. \quad (45)$$

Various special cases of this photon distribution function were investigated also in [16,18]. Confining ourselves to the case of $\langle p \rangle = \langle q \rangle = 0$ and taking into account eq. (4), we can simplify (41) as follows,

$$\mathcal{P}_n = \left(d + \frac{T}{2} + \frac{1}{4}\right)^{-\frac{1}{2}} \left(\frac{4d + 1 - 2T}{4d + 1 + 2T}\right)^{\frac{n}{2}} P_n \left(\frac{4d - 1}{[(4d + 1)^2 - 4T^2]^{\frac{1}{2}}}\right). \quad (46)$$

For *pure quantum states*, when $d = 1/4$, we have an *oscillating* photon distribution function [5,19,20]:

$$\mathcal{P}_{2m}^{(pure)} = \left(\frac{2}{1 + T}\right)^{\frac{1}{2}} \frac{(2m)!}{(2^m m!)^2} \left(\frac{T - 1}{T + 1}\right)^m, \quad \mathcal{P}_{2m+1}^{(pure)} = 0, \quad (47)$$

To see the “smoothing” of these oscillations for *mixed* quantum states we use eq. (25). Since in the case under study

$$r = \frac{1 - 4d}{2\sqrt{T^2 - 4d}} \leq 0,$$

we choose the values of T and d in the domain

$$T - \frac{1}{2} > 2d \geq \frac{1}{2},$$

to ensure the inequality $|r| < 1$. Then we have for $n \gg 1$

$$\mathcal{P}_n \approx \frac{[2\sqrt{T^2 - 4d} + 4d - 1]^{n+\frac{1}{2}} + (-1)^n [2\sqrt{T^2 - 4d} - 4d + 1]^{n+\frac{1}{2}}}{\sqrt{\pi n}(1 + 2T + 4d)^{n+\frac{1}{2}}(T^2 - 4d)^{\frac{1}{4}}}. \quad (48)$$

For $d = 1/4$ this expression coincides with the Stirling asymptotics of formula (47). Eq. (48) shows that the oscillations of the photon distribution function disappear provided

$$n \frac{4d-1}{2\sqrt{T^2-4d}} \geq 1.$$

For highly squeezed mixed states, when $T^2 \gg 4d \gg 1$, this criterion assumes the form $2nd/T \geq 1$.

3. For a quantum harmonic oscillator with a time-dependent frequency the transition probability from the n -th to the m -th stationary level can be expressed as [1-3,7,8]

$$W_n^m = \frac{W_0^0}{n!m!} \left| H_{nm}^{\{\mathbf{R}\}}(x_1, x_2) \right|^2, \quad (49)$$

with matrix $\mathbf{R}$ of the form

$$\mathbf{R} = -\frac{1}{\xi} \begin{pmatrix} -\eta & 1 \\ 1 & \eta^* \end{pmatrix}, \quad |\xi|^2 - |\eta|^2 = 1,$$

and

$$W_0^0 = |\xi|^{-1}.$$

The explicit form of the parameters ξ and η is determined by the solutions of the classical equation

$$\ddot{\epsilon} + \omega^2(t)\epsilon = 0,$$

moreover, the ratio $|\eta/\xi|^2$ may be interpreted as a reflection coefficient from some “effective potential barrier” $\omega^2(t)$, provided a “transmitted wave” is $\exp(i\omega t)$ for $t \rightarrow -\infty$. The parameters x_1, x_2 are expressed through the convolution of the function $\epsilon(t)$ with the external time-dependent force. If this force is absent, $x_1 = x_2 = 0$. Confining ourselves with this simplest case we can rewrite eq. (49), due to eqs. (5) and (6), in the following equivalent forms (remind that m and n must be either both even or both odd):

$$W_n^m = \frac{n!m!}{\left[\left(\frac{n+m}{2}\right)!\right]^2 |\xi|} \left| \frac{\eta}{2\xi} \right|^{|n-m|} \left[P_{\mu_{nm}}^{\left(\frac{|n-m|}{2}, \frac{|n-m|}{2}\right)} \left(\frac{1}{|\xi|} \right) \right]^2, \quad (50)$$

$$W_n^m = \frac{\mu_{nm}!}{\nu_{nm}!} \left[\frac{|n-m|!}{\left(\frac{|n-m|}{2}\right)!} \right]^2 \frac{1}{|\xi|} \left| \frac{\eta}{2\xi} \right|^{|n-m|} \left[C_{\mu_{nm}}^{\frac{|n-m|+1}{2}} \left(\frac{1}{|\xi|} \right) \right]^2. \quad (51)$$

This is just the case when the argument of the Jacobi and Gegenbauer polynomials belongs to the interval $[-1, 1]$. Therefore the asymptotics for $\mu_{nm} \gg 1$, $|n-m| \sim \mathcal{O}(1)$ is given by eq. (26):

$$W_n^m \approx \frac{\arctan|\eta|}{|\eta|} \left[J_{\frac{|n-m|}{2}}(\mathcal{N}_{nm} \arctan|\eta|) \right]^2. \quad (52)$$

If $\mathcal{N}_{nm} \arctan|\eta| \gg 1$, the asymptotics of the Bessel function yields

$$W_n^m \approx \frac{2}{\pi \mathcal{N}_{nm} |\eta|} \left[\cos \left(\mathcal{N}_{nm} \arctan|\eta| - \frac{\pi}{4} (|n-m| + 1) \right) \right]^2. \quad (53)$$

If $|n - m| \gg \mu_{nm} \sim \mathcal{O}(1)$, then eq. (28) leads to the asymptotics

$$W_n^m \approx \sqrt{\frac{2}{\pi \nu_{nm}}} \frac{|\eta|^{n-m}}{2^{\mu_{nm}} \mu_{nm}! |\xi|^{n-m+1}} \left[H_{\mu_{nm}} \left(\sqrt{\frac{|n-m|+1}{2|\xi|^2}} \right) \right]^2. \quad (54)$$

For $|\xi| \sim \mathcal{O}(1)$ this formula can be simplified as follows,

$$W_n^m \approx \sqrt{\frac{2}{\pi \nu_{nm}}} \frac{\nu_{nm}^{\mu_{nm}} |\eta|^{n-m}}{\mu_{nm}! |\xi|^{n+m+1}}. \quad (55)$$

Two other special cases correspond to small and large values of the “reflection coefficient $|\eta/\xi|^2$ ”. If $|\eta| \rightarrow 0$ (this limit corresponds to an adiabatic change of the frequency $\omega(t)$, when $|\xi| \rightarrow 1$), we get from eq. (51)

$$W_n^m \approx \frac{\nu_{nm}!}{\mu_{nm}! \left[\left(\frac{|n-m|}{2} \right)! \right]^2} \left| \frac{\eta}{2} \right|^{n-m}. \quad (56)$$

For large values of n and m this expression coincides with the limits of eqs. (52) and (55) for $|\eta| \rightarrow 0$, if one takes into account formula (22).

An interesting phenomenon is observed when $|\xi| \approx |\eta| \gg 1$ (more precisely, when $|\xi| \gg \nu_{nm}$). Then for even values of n and m we get from eq. (51)

$$W_{2k}^{2l} \approx \frac{1}{|\xi|} \frac{(2k-1)!!}{(2k)!!} \frac{(2l-1)!!}{(2l)!!}, \quad (57)$$

whereas the transition probability between the neighbouring odd levels is much less:

$$W_{2k+1}^{2l+1} \approx \left[\frac{2\nu_{kl}+1}{|\xi|} \right]^2 W_{2k}^{2l}. \quad (58)$$

For instance,

$$W_0^0 = |\xi|^{-1}, \quad W_1^1 = |\xi|^{-3}, \quad W_2^2 \approx \frac{1}{4} |\xi|^{-1}, \quad W_3^3 \approx \frac{9}{4} |\xi|^{-3}, \dots$$

So, for $|\xi| \gg \nu_{nm}$ the transition probabilities oscillate like the photon distribution function in eq. (47). Such a coincidence is not accidental, due to the relation [3]

$$|\xi| = \sqrt{\mathcal{E} + \frac{1}{2}},$$

$\mathcal{E}$ being the normalized energy of quantum fluctuations in the final Gaussian state which has originated from the initial ground state with $\mathcal{E}_{\text{in}} = \frac{1}{2}$. It is well known [3,21] that Gaussian states with large energy of fluctuations are highly squeezed, and a manifestation of squeezing is just the oscillating structure of various distributions related to the number of quanta.

7 Acknowledgement

One of us (V.I.M.) thanks INFN and University of Napoli "Federico II" for the hospitality.

References

- [1] I.A.Malkin, V.I.Man'ko, and D.A.Trifonov, J. Math. Phys. **14**, 576 (1973).
- [2] I.A.Malkin and V.I.Man'ko, *Dynamical Symmetries and Coherent States of Quantum Systems* (Nauka, Moscow, 1979; in Russian).
- [3] V.V.Dodonov and V.I.Man'ko, *Invariants and Evolution of Nonstationary Quantum Systems*, Proceedings of Lebedev Physics Institute, v.**183**, edited by M.A.Markov (English translation: Nova Science Publishers, Commack, N.Y., 1989).
- [4] M.Kauderer, J. Math. Phys. **34**, 4221 (1993).
- [5] A.Vourdas and R.M.Weiner, Phys. Rev. **36**, 5866 (1987).
- [6] G.Adam, Phys.Lett.A **171**, 66 (1992).
- [7] I.A.Malkin, V.I.Man'ko, and D.A.Trifonov, Phys. Rev. D **2**, 1371 (1970).
- [8] V.V.Dodonov, I.A.Malkin, and V.I.Man'ko, Physica **59**, 241 (1972).
- [9] V.V.Dodonov, V.I.Man'ko, and V.V.Semjonov, Nuovo Cimento B **83**, 145 (1984).
- [10] V.V.Dodonov, V.I.Man'ko, and I.N.Prokopenya, *Asymptotic Density Matrix Elements of the Canonically Transformed Oscillator Hamiltonian in the Fock Basis*, Preprint No. 30 (Lebedev Physics Institute, Moscow, 1985).
- [11] A.Vourdas, Phys.Rev.A **34**, 3466 (1986).
- [12] *Bateman Manuscript Project: Higher Transcendental Functions*, edited by A.Erdélyi (McGraw-Hill, New York, 1953).
- [13] G.Szegö, *Orthogonal Polynomials* (American Math. Society, New York, 1959).
- [14] F.W.J.Olver, *Asymptotics and Special Functions* (Academic Press, New York, 1974).
- [15] *Handbook of Mathematical Functions*, edited by M.Abramowitz and I.A.Stegun (National Bureau of Standards, Washington, D.C., 1964).
- [16] S.Chaturvedi and V.Srinivasan, Phys.Rev.A **40**, 6095 (1989).
- [17] V.V.Dodonov, O.V.Man'ko, and V.I.Man'ko, "Photon distribution for one-mode mixed light with generic gaussian Wigner function", University of Napoli preprint INFN-NA-IV-93/35, DSF-T-93/35(1993) (submitted to Phys. Rev. A).
- [18] P.Marian and T.A.Marian, Phys.Rev.A **47**, 4474 (1993).
- [19] W.Schleich and J.A.Wheeler, J.Opt.Soc.Am.B **4**, 1715 (1987).
- [20] V.V.Dodonov, A.B.Klimov, and V.I.Man'ko, Phys.Lett.A **134**, 211 (1989).
- [21] V.V.Dodonov, A.B.Klimov, and V.I.Man'ko, in *Squeezed and Correlated States of Quantum Systems*, Proceedings of Lebedev Physics Institute, v. **205**, edited by M.A.Markov (Nova Science, Commack, 1993).